

# New Features of HYCOM

Alan J. Wallcraft

Naval Research Laboratory

15th Layered Ocean Model Workshop

June 2, 2009

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE <b>02 JUN 2009</b>		2. REPORT TYPE		3. DATES COVERED <b>00-00-2009 to 00-00-2009</b>	
4. TITLE AND SUBTITLE <b>New Features of HYCOM</b>				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) <b>Naval Research Laboratory,Stennis Space Center,MS,39529</b>				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT <b>Approved for public release; distribution unlimited</b>					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT <b>Same as Report (SAR)</b>	18. NUMBER OF PAGES <b>21</b>	19a. NAME OF RESPONSIBLE PERSON
a. REPORT <b>unclassified</b>	b. ABSTRACT <b>unclassified</b>	c. THIS PAGE <b>unclassified</b>			

## **HYCOM 2.2 (I)**

- Maintain all features of HYCOM 2.1
  - Orthogonal curvilinear grids
  - Can emulate Z or Sigma or Sigma-Z models
    - ◇ It is “Arbitrary Lagrangian-Eulerian”, see:  
Adcroft and Hallberg, O. Modelling 11 224-233.
  - Explicit support for 1-D and 2-D domains
  - KPP or Kraus-Turner or Mellor-Yamada 2.5 or Price-Weller-Pinkel
  - Rivers as bogused surface precipitation
  - Multiple tracers
  - Off-line one-way nesting
  - Scalability via OpenMP or MPI or both
    - ◇ Bit-for-bit multi-cpu reproducibility
- Special halo exchange for tripole global grid
  - Arctic dipole patch on standard Mercator globe
  - Logically rectangular domain
    - ◇ Two halves of top edge “fold” together
    - ◇ V-velocity changes sign across the fold

## **HYCOM 2.2 (IIa)**


- Alternative LeapFrog barotropic time splitting
  - Provided by SHOM
  - Twice as expensive as standard scheme
    - ◇ Still a small fraction of total run time
  - Significantly more stable
  - May allow 2x longer baroclinic time step
- Alternative scalar advection techniques
  - Provided by Mohamed Iskandarani
  - Donor Cell, FCT (2nd and 4th order), MPDATA
  - FCT2 replaces MPDATA as standard scheme
- Mixed layer changes
  - GISS mixed layer model
    - ◇ Provided by Armando Howard
  - KPP bottom boundary layer
    - ◇ Provided by George Halliwell
  - KPP tuning

## **HYCOM 2.2 (IIb)**


- Initial vertical coordinate changes
  - Must always use PCM for isopycnal layers
  - Vertical remapping used PLM for fixed coordinate layers
  - Thin deep iso-pycnal layers
  - Stability from locally referenced potential density
  - Spatially varying layer target densities
    - ◇ Different isopycnal layers in semi-enclosed seas
- Recent vertical coordinate changes
  - hybrlx only active below “fixed coordinate” surface layers
  - Major re-write of HYBGEN by George Halliwell and Alan Wallcraft
    - ◇ Must always use PCM for isopycnal layers
    - ◇ Vertical remapping uses PLM or PPM or WENO-like PPM (Laurent Debreu) for fixed and non-isopycnal coordinate layers
    - ◇ More layers are identified as non-isopycnal
    - ◇ Updated logic for two layers (one too dense, other too light)

# ANNUAL MEAN EQUATORIAL PACIFIC GLBt0.72 HYCOM VERSION 2.2.03

**u-velocity zonal sec. 0.00n mean: 4.004- 5.004 [06.0H]**




**temperature zonal sec. 0.00n mean: 4.004- 5.004 [06.0H]**




# ANNUAL MEAN EQUATORIAL PACIFIC

## GLBt0.72 HYCOM VERSION 2.2.18

**u-velocity zonal sec. 0.00n mean: 4.004- 5.004 [22.3H]**



**temperature zonal sec. 0.00n mean: 4.004- 5.004 [22.3H]**



## HYCOM 2.2 (IIc)

- Equation of state that is quadratic in salinity
  - HYCOM must “invert” the equation of state
    - ◇ tofsig(r,s) and sofsig(r,t)
  - Traditional version is cubic in T and linear in S
    - ◇ Finding the root of a cubic is expensive, but exact
    - ◇ Linear in S is not accurate at low salinity
  - Optional version is cubic in T and quadratic in S
    - ◇ Coefficients provided by Shan Sun
    - ◇ More accurate at low salinity
      - Rivers, Black Sea, Caspian Sea
    - ◇ Not much more expensive
- Pade equation of state
  - Optional Pade version:  $P_{22}/Q_{11}$ 
    - ◇  $P_{22}$  is quadratic in T and S
    - ◇  $Q_{11}$  is linear in T and S
  - Developed at NCEP
    - ◇ Only Sigma0 at present
  - More accurate at low salinity




## **HYCOM 2.2 (IId)**

- Improved thermobaricity
  - Use exact formula for the thermobaricity
    - ◇ Eqn. 3 from Sun et al. 1999
  - No single reference state is appropriate for the global ocean
    - ◇ Hallberg, Ocean Modelling, 8, 279-300
  - Use a linear combination of pressure gradients from two out of three reference states
    - ◇ Atlantic (3°C, 35.0 psu)
    - ◇ Arctic/Antarctic (0°C, 34.5 psu)
    - ◇ Mediterranean (13°C, 38.5 psu)
  - Most locations use just one reference state
    - ◇ Linear combinations allow smooth transition between states
      - Do this in shallow water if possible
  - In deep water, constrain the T&S used for thermobaricity to be close to the reference state


## HYCOM 2.2 (IIIa)

- Atmospheric forcing changes
  - Option to input ustar fields
    - ◇ Best option for monthly forcing
    - ◇ Otherwise calculated from wind stress or speed
  - Can relax to observed SST fields
  - Improved COARE 3.0 bulk exchange coefficients
  - Black-body correction to longwave flux
  - Climatological heat flux offset,  $\overline{Q}_c$ 
    - $$Q = (Q_{sw} - Q_{lw}) + (Q_l + Q_s) + \overline{Q}_c$$
    - ◇  $\overline{Q}_c$  is constant in time
      - Typically based on the model's climatological SST error, times (say)  $-45 \text{ W m}^{-2} / ^\circ\text{C}$
  - Analytic diurnal heat flux cycle
    - ◇ Need hourly  $Q_{sw}$  for good cycle
      - Have 3 or 6 hourly (snapshots or averages)
    - ◇ Input daily-running average  $Q_{sw}$
    - ◇ Apply multiplicative correction:
      - $\text{clear-sky\_now} / \text{clear-sky\_daily-average}$


# 1-D HYCOM at OCEAN STATION PAPA NOGAPS FORCING



SST in 2007 at 145W,50E



ShortWave Flux in 2007 at 145W,50E



## HYCOM 2.2 (IIIb)

- Improved support for rivers
  - Still bogused surface precipitation
  - High frequency inter-annual river flow allowed
    - ◇ Add it to atmospheric precip, off-line
    - ◇ Instead of monthly climatology, or in-addition to it (flow anomalies)
  - Better control of low salinity profiles
  - Option for mass (vs salinity) flux
  - Equation of state that is quadratic in salinity
- Tidal forcing
  - Provided by NCEP
  - Body forcing and open boundary forcing
  - Boundary forcing currently for “Flather” ports
    - ◇ v2.3: Extend it to Browning-Kreiss nesting
  - Body forcing for 8 largest components
  - SAL treated as a fraction of non-steric SSH
  - Tidal drag based on bottom roughness
    - ◇ Applied to near-bottom tidal velocity
      - Use a lagged 25-hour average as the non-tidal velocity

## **HYCOM 2.2 (IIIc)**

- New diagnostics within HYCOM
  - Time-averaged fields (in archive files)
    - ◇ Identical to off-line mean archives
    - ◇ Daily 25-hour average for tides
    - ◇ No in-line capability to capture variability
    - ◇ Instantaneous archives still available
  - Sub-region archive files
    - ◇ Example: hourly 3-D from Global 1/12°
      - 3-4 small regions only
    - ◇ One file per involved MPI task (entire tile)
    - ◇ Reconstruct standard regional archive files off-line
    - ◇ Instantaneous archives still available
  - Synthetic instrumentation
    - ◇ Provided by George Halliwell
    - ◇ 3-D particle tracking
    - ◇ surface and constant depth drifters
    - ◇ isopycnic drifters
    - ◇ fixed instruments and moorings

## **HYCOM 2.2 (IIIId)**

- I/O optimizations
  - Typically, all I/O is from a single MPI task
  - I/O can be a bottleneck when running on many processors
    - ◇ MPI-2 I/O option
      - Do I/O from 1st MPI task in each row of tiles
    - ◇ Faster code for “endian” conversion
      - HYCOM files are always big-endian
      - Intel and AMD are little-endian
      - Do the conversion in parallel
  - Sub-region archive files
    - ◇ One file per involved MPI task
    - ◇ Can be much faster than writing a full archive
  - Remove density from restart and archive files
    - ◇ Less I/O, smaller files
    - ◇ Must track which equation of state is used

## HYCOM 2.2 AND SEA ICE

- Finer control over energy loan ice model
  - Melting point can be linear in salinity
  - Set ice minimum and maximum thickness
  - Set ice vertical temperature gradient
    - ◇ Or get ice surface temperature from  $T_a$
  - Made compatible with coupled sea-ice approach
- Two-way coupling to LANL's CICE sea ice model
  - HYCOM exports:
    - ◇ SST, SSS, SSH
    - ◇ Surface Currents
    - ◇ Available Freeze/Melt Heat Flux
  - CICE exports:
    - ◇ Ice Concentration
    - ◇ Ice-Ocean Stress
    - ◇ Actual Freeze/Melt Heat/Salt/Mass Flux
    - ◇ Solar Radiation at Ice Base
  - Coupling via the Earth System Modeling Framework
    - ◇ Currently for non-global domains only
    - ◇ v2.3: ESMF version 4 and global domains



## **HYCOM 2.2 AND CCSM**

- Community Climate System Model  
**<http://www.ccsm.ucar.edu/>**
  - Fully-coupled, global climate model
  - Sea-Ice: CICE
  - Ocean: POP
- HYCOM can be used in place of POP in CCSM3
- Uses the standard HYCOM source code
- Subdirectory CCSM3 used to hold and build the CCSM3 version
  - Some source code files are specific to CCSM3
  - HYCOM ".f" files are renamed ".F" to simplify CCMS3 integration
  - Macro USE\_CCSM3 for CCSM3-specific code

## **HYCOM 2.2 (IV)**


- Climatological nesting now allowed
  - Start from monthly mean outer model archive files
  - Allows nested runs longer than the outer run
    - ◇ But with less accurate boundary state
  - Probably only suitable for regional nests
- Nesting no longer requires co-located grids
  - General archive to archive horizontal interpolation (curvilinear)
- Hybrid to fixed vertical grid remapper
  - Allows fixed-coordinate nests inside hybrid coordinate outer domains
    - ◇ HYCOM to (fixed-grid) HYCOM
    - ◇ HYCOM to NCOM

## **HYCOM 2.2 (V)**

- Enhanced hycomproc and fieldproc
  - NCAR-graphics based
  - Many more color palette options
    - ◇ Can read in an arbitrary palette
  - Mark locations, and draw tracks, on plot
  - Plot diffusion coefficients and tracers (hycomproc)
  - Overlay vector and line-contours (fieldproc)
    - ◇ Vectors can optionally follow streamlines
- Added fieldcell
  - Like fieldproc, but for cell-array (vs contouring)
    - ◇ Mark locations and draw tracks
    - ◇ Overlay line-contours
  - Uses NCAR's map projections
  - Typically much faster than fieldproc, but can leave unfilled cells
  - Option to increase resolution of input (bi-linear interpolation)

# EXAMPLE OF FIELDPROC SSH, SURFACE CURRENTS, AND BATHYMETRY

**FSD (cm) and V.1 - Jan Year 20**



## **HYCOM 2.2 (VI)**

- Diagnostic fields to netCDF and other file formats
  - Archive fields in layer space
    - ◇ On p-grid (interpolated velocity)
  - 3-D archive fields interpolated to z-space
    - ◇ On p-grid, or
    - ◇ Sampled at stations or along arbitrary tracks
  - 3-D archive fields sampled on iso-therms
  - Meridional stream-function from (mean) 3-D archive
    - ◇ In logical array space (rectilinear grids)
    - ◇ Binned to latitude bands (curvilinear grids)
  - Atmospheric forcing input fields
    - ◇ Time axis depends on “.b” file format
    - ◇ Any “.a” file with the right “.b” file structure can be converted to netCDF
  - Fields binned into lon-lat cells

## **CANDIDATE FEATURES FOR HYCOM 2.3**

- Wind drag coefficient based on model SST
- Regional tides and Browning-Kreiss nesting
- Wetting and Drying
- Fully region-independent
  - Compile once, run on any region and any number of processors
    - ◇ Run-time memory allocation
    - ◇ Might reduce performance (fewer compiler optimizations available)
  - Needed for full ESMF compliance
- Enhanced support for ESMF
  - HYCOM+CICE on global tripole grid